

The Abnormal Pap Smear

Laurie Turenne-Kolpan, Maj,
USAF, MC

Objectives

- Review the epidemiology of cervical cancer
- Review risk factors for cervical cancer
- Review screening recommendations
- Discuss the reporting of pap smear results using the Bethesda System
- Determine the appropriate management of abnormal pap smear results

Introduction

- The Papanicolaou (Pap) smear has been in use since 1941
- It is the single most effective cancer screening test to date
- Sensitivity is 30-89%

Epidemiology

- 3.5 million women have abnormal pap results in the US annually (about 7% of total paps performed)
- 4.4% of pap results are ASCUS
- In the US, 13,000 new cases of invasive cervical cancer per year
- 50% of women in the US with cervical cancer have never been screened
- 5-year survival rate for local disease is 92%
- 5-year survival for distant metastasis is 13%

Risk Factors for Cervical Cancer

- Early age at first intercourse (age <16)
- Multiple sexual partners
- Presence of STD, particularly HPV
- Immunosuppression
- SMOKING!!!

The Human Papillomavirus (HPV)

- HPV is the leading etiologic agent in development of dysplasia
- HPV DNA is found in 95-100% of invasive cervical cancer (ICC) and 75-95% of high grade lesions (CIN II or III)
- Peak prevalence is 40% occurring between the ages of 20 and 29
- Condoms are not as protective against transmission of HPV as other STDs

(HPV – continued)

- High risk types –
16,18,31,33,35,39,45,51,52,56,58,59,68,
73,82
- Low risk types – 6,11,42,43,44, 54,
61,70,72,81
- Types 16 and 18 most prevalent in ICC
- Persistence of HPV infection is a key
factor in progression to cancer

Screening - When to Begin?

- New guidelines! - ACOG, American Cancer Society (ACS), and US Preventive Services Task Force (USPSTF) all recommend screening starting 3 years after onset of sexual activity, or at the age of 21

Screening – When to end?

- ACOG – No set upper age limit
- USPSTF – 70, if have had 3 consecutive negative tests, and no abnormalities in the last 10 years
- ACS – 65, if they are not at high risk and have adequate previous screenings
- After total hysterectomy for benign disease (ACOG adds also if no h/o CIN II or III)

How Often to Screen?

- **ACOG** – annually for all women under age 30 regardless of method (conventional vs. liquid)
 - Over age 30 can space to every 2-3 years IF
 - No history of CIN II or III
 - Not immunocompromised
 - Not HIV positive
 - Had no DES exposure
 - Have had 3 consecutive negative screens

How Often (Cont)?

- **ACS** - Annual with conventional cytology, or every 2 years with liquid based cytology
 - **After age 30, may increase interval to every 2-3 years IF**
 - **Had 3 consecutive negative screens**
 - **Not high risk**
 - **Not immunocompromised**
- **USPSTF** - At least every 3 years

Which test to use?

- Conventional pap smear is about 60-80% sensitive for detecting a high grade lesion
- Newer tests including liquid-based paps and computerized re-screening have been shown in some studies to have a higher sensitivity, but lower specificity
- No prospective trials comparing conventional with new technologies in prevention of ICC or cost-effectiveness

Bethesda 2001

- Specimen type (conventional, liquid-based, etc)
- Specimen Adequacy
 - Satisfactory (describes presence or absence of endocervical/ TZ component and other quality indicators such as obscuring blood or inflammation)
 - Unsatisfactory (reason specified)
- General Categorization
 - Negative for intraepithelial lesion or malignancy
 - Epithelial Cell Abnormality
 - Other (endometrial cells in a woman over 40)

Absent Endocervical Component

- Expert task force has recommended that repeat cytology can be performed in 12 months. Should repeat in 6 months IF:
 - \geq ASCUS pap without 3 consecutive normals
 - H/O atypical glandular cells of unknown origin
 - + Hi risk HPV in the last 12 months
 - Inability to visualize or sample the endocervical canal
 - Immunosuppression
 - Non-compliant patient

Negative for Intraepithelial

- **Organisms**
 - *Trichomonas vaginalis*
 - Fungal organisms
 - Shift in flora suggestive of BV
 - Cellular changes c/w Herpes
 - *Actinomyces* spp.
- **Other non-neoplastic findings (optional)**
 - Reactive cellular changes
 - Glandular cells post hysterectomy
 - Atrophy


Bethesda 2001 - Epithelial cell abnormalities - Squamous

- Atypical squamous cells
 - ASC-US (of undermined significance)
 - ASC-H (cannot exclude HSIL)
- Low grade squamous intraepithelial lesion (LSIL) – mild dysplasia/ CIN I
- High grade squamous intraepithelial lesion (HSIL) – moderate and severe dysplasia/ CIN II/III
- Squamous cell carcinoma


Bethesda 2001 - Epithelial cell abnormalities -

- Atypical **Glandular**
 - Endocervical (NOS)
 - Endometrial (NOS)
 - Glandular (NOS)
- Atypical
 - Endocervical, favor neoplastic
 - Glandular, favor neoplastic
- Endocervical adenocarcinoma *in situ*
- Adenocarcinoma - endocervical, endometrial, extrauterine, NOS


Management of Women with Atypical Squamous Cells of Undetermined Significance (ASC-US)




Management of Women with Atypical Squamous Cells of Undetermined Significance (ASC-US) In Special Circumstances



Management of Women with Atypical Squamous Cells: Cannot Exclude High-grade SIL (ASC - H)




Management of Women with Low-grade Squamous Intraepithelial Lesions (LSIL) *




* Management options may vary if the woman is pregnant, postmenopausal, or an adolescent - (see text)

Management of Women with Low-grade Squamous Intraepithelial Lesions In Special Circumstances




* For low-risk, post-menopausal women with a history of negative screening

Management of Women with Low-grade Squamous Intraepithelial Lesions In Special Circumstances




Management of Women with High-grade Squamous Intraepithelial Lesions (HSIL) *



^ Includes referral cytology, colposcopic findings, and all biopsies

* Management options may vary if the woman is pregnant, postmenopausal, or an adolescent

Management of Women with Atypical Glandular Cells (AGC)



Summary

- Pap smear testing – single most effective cancer screening test to date
- Risk factors: HPV and other STDs, early intercourse, multiple partners, immunosuppression, smoking
- Begin screening 3 yrs after onset of sexual activity, or at age 21
- Persistence of HPV is key factor in developing high grade dysplasia/ICC
- HPV typing can be used to triage ASCUS paps
- Remember ASCCP clinical guidelines for management of cytological abnormalities

References

- Wright, Jr TC, Cox JT, Massad LS, Twiggs, LB, Wilkinson, EJ. 2001 Consensus Guidelines for the Management of Women with Cervical Cytological Abnormalities. JAMA 2002; 287: 2120-2129.
- Holowaty P, Miller AB, Rohan T, To T. Natural History of Dysplasia of the Uterine Cervix. Obstetrical & Gynecological Survey 1999; 54: 375-377.
- Sawaya GF, Brown AD, Washington AE, Garber AM. Current Approaches to Cervical-Cancer Screening. The New England Journal of Medicine 2001; 344: 1603-1607.
- Grimm KJ, Meadows SE, Ruplinger JM. Who should have colposcopy? The Journal of Family Practice 2003; 52: 64-66.
- U.S. Preventive Services Task Force. Screening for Cervical Cancer: Recommendations and Rationale. American Family Physician 2003; 8: 1759-1769.
- Altekruze SF, Lacey JV, Jr, Brinton LA, Gravitt, PE, Silverberg SG, Barnes WA, Jr, Greenberg MD, Hadjimichael OC, McGowan L, Mortel R, Schwartz PE, Hildesheim A. Comparison of human Papillomavirus genotypes, sexual and reproductive risk factors of cervical adenocarcinoma and squamous cell carcinoma. American Journal of Obstetrics and Gynecology 2003; 188: 657-663 (5)
- Murthy NS, Mathew A. Risk factors for pre-cancerous lesions of the cervix. European Journal of Cancer Prevention 2000; 9:
- Goodman A, Holschneider C. Management of the abnormal papanicolaou smear. Up to Date 2004.
- Solomon D et al. The 2001 Bethesda System. JAMA 2002: 287 (16)